

Accepted Manuscript

Endoscopically-Assisted Surgical Expansion (EASE) for the Treatment of Obstructive Sleep Apnea

Kasey Li, DDS, MD, FACS, Stacey Quo, DDS, MS, Christian Guilleminault, MD



PII: S1389-9457(18)30391-5

DOI: [10.1016/j.sleep.2018.09.008](https://doi.org/10.1016/j.sleep.2018.09.008)

Reference: SLEEP 3820

To appear in: *Sleep Medicine*

Received Date: 11 July 2018

Revised Date: 8 September 2018

Accepted Date: 12 September 2018

Please cite this article as: Li K, Quo S, Guilleminault C, Endoscopically-Assisted Surgical Expansion (EASE) for the Treatment of Obstructive Sleep Apnea, *Sleep Medicine* (2018), doi: <https://doi.org/10.1016/j.sleep.2018.09.008>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Endoscopically-Assisted Surgical Expansion (EASE) for the Treatment of Obstructive Sleep Apnea

Kasey Li, DDS, MD, FACS^{a, *}, Stacey Quo, DDS, MS^b, Christian Guillemineault, MD^c

^aSleep Apnea Surgery Center, East Palo Alto, CA, USA

^bSchool of Dentistry, University of California at San Francisco, San Francisco, CA, USA

^cSleep Medicine Division, Stanford University School of Medicine, Stanford, CA, USA

*Corresponding Author: Kasey Li, DDS, MD, FACS

E-mail: drli@sleepapneasurgery.com

1900 University Avenue

Suite 105

East Palo Alto, CA 94303

USA

Abstract

Objective: The aim of this retrospective study was to evaluate the results of an outpatient surgical procedure named endoscopically-assisted surgical expansion (EASE) in expanding the maxilla to treat obstructive sleep apnea (OSA) in adolescent and adults.

Methods: Thirty-three patients (18 males), aged 15-61 years, underwent EASE of the maxilla. All patients completed pre- and post-operative clinical evaluations, polysomnography, questionnaires (Epworth Sleepiness Scale [ESS] and Nasal Obstruction Septoplasty Questionnaire [NOSE]) and cone beam computed tomography (CBCT).

Results: With EASE, the overall apnea hypopnea index (AHI) improved from 31.6 ± 11.3 to 10.1 ± 6.3 . The oxygen desaturation index (ODI) improved from 11.8 ± 9.6 to 1.8 ± 3.7 , with reduction of ESS scores from 13.4 ± 4.0 to 6.7 ± 3.1 . Nasal breathing improved as demonstrated by reduction of the NOSE scores from 57.8 ± 12.9 to 15.6 ± 5.7 . Expansion of the airway from widening of the nasal floor was consistently evident on all postoperative CBCT; the anterior nasal floor expanded 4.9 ± 1.2 mm, posterior nasal floor expanded 5.6 ± 1.2 mm, and the dental diastema created was 2.3 ± 0.8 mm. Mean operative time was 54.0 ± 6.0 minutes. All patients with mild to moderate OSA were discharged the same day; patients with severe OSA were observed overnight. All patients returned to school or work and regular activities within 3 days.

Conclusions: EASE is an outpatient procedure that improves nasal breathing and OSA by widening the nasal floor in adolescents and adults. Compared to current surgical approaches for maxillary expansion, EASE is considerably less invasive and consistently achieves enlargement of the airway with minimal complications.

Keywords:

Maxillary expansion, palatal expansion, SARPE, EASE, OSA, airway, SARME

ACCEPTED MANUSCRIPT

1. Introduction

Since the first report of maxillary expansion for the treatment of obstructive sleep apnea (OSA) by Cistulli et al. in 1998 (1), its efficacy in expanding the airway and improving OSA has been validated repeatedly (2-5). However, the majority of studies regarding maxillary expansion to treat OSA have occurred in the pediatric population. This is because maxillary expansion is performed easily in children via use of an orthodontic expander as part of a routine, non-invasive procedure.

As the maxilla matures during puberty, ossification of the midpalatal suture occurs with resultant posterior-to-anterior formation of mineralized bridges (6). In adults and adolescents, maxillary expansion is more challenging due to skull maturation, which results in increased resistance to suture separation (7). Therefore, surgically-assisted rapid palatal expansion (SARPE) is typically needed to facilitate widening of the maxilla in non-growing patients (8-10). However, SARPE is an invasive procedure associated with potential complications, which include significant hemorrhage, excessive lacrimation, loss of bone and teeth, significant pain and numbness (11, 12), as well as aesthetic changes including nasal base widening and lip shortening (13-15). In addition, SARPE is associated with a prolonged recovery due to residual pain and swelling. Most importantly, SARPE may not achieve the surgical goal of expanding the airway, especially in the posterior nasal aspect (Fig. 1).




Fig. 1. (a) Frontal view of CBCT demonstrating maxillary widening between the roots of central incisors. (b) Frontal view showing the dental diastema being closed. (c) Palatal view demonstrating that the nasal floor was widened in a tapered fashion (black arrows), but the posterior region remained unchanged (white arrow). (d) Palatal view showing that further expansion led to greater anterior widening (black arrows) but failed to open the posterior nasal airway (white arrow).

In 2004, Guilleminault and Li reported their initial experiences using SARPE to surgically expand the maxilla and using mandibular widening to treat OSA (16). To reduce the invasiveness, minimize risks and improve outcomes of maxillary expansion, the surgical technique has evolved over the past 15 years. The aim of this retrospective study was to assess the results of using endoscopically-assisted surgical expansion (EASE) as a

significantly less invasive approach of maxillary widening surgery for the treatment of OSA.

2. Materials and Methods

2.1. Subjects

This study retrospectively reviewed patients (aged 15 or older) who underwent EASE of the maxilla to treat OSA. Data evaluated included clinical and operating room records, polysomnography (PSG) records, Epworth Sleepiness Scale (ESS) questionnaires, Nasal Obstruction Septoplasty Effectiveness (NOSE) questionnaires and cone beam computed tomography (CBCT) results.

2.2. Surgical Procedure: Endoscopically-assisted surgical expansion (EASE)

The same surgical procedure, specifically EASE, was performed in all patients. Under general anesthesia, a transpalatal distractor (TPD, KLS Martin Group, Jacksonville, FL) was inserted onto the palate at the region of the first molar. The TPD was activated such that the footplates fully engaged the bone, and each footplate was stabilized with a screw. A stab incision in the posterior tuberosity was made, and the pterygomaxillary suture was identified using a periosteal elevator. Gentle pterygomaxillary separation was achieved with a piezoelectric blade (DePuy Synthes, Switzerland). During separation of the pterygomaxillary suture, a finger was placed intraorally to palpate separation of the suture while avoiding injury to the intraoral mucosa. Using a nasal endoscope for visualization, midpalatal osteotomy was performed with a piezoelectric blade. The midpalatal osteotomy was initiated from the posterior nasal spine (PNS) at the junction between the nasal septum and nasal floor with the blade angling towards the midline (Fig. 2). The entire osteotomy was performed with the piezoelectric blade cutting through the nasal mucosa and bone while taking care to avoid injuring palatal mucosa. The osteotomy was carried anteriorly to within 2-3 mm of the anterior nasal spine (ANS). The

ANS and bone in between the roots of the central incisors was not disturbed. The midpalatal osteotomy was performed bilaterally to ensure symmetrical separation of the midpalatal suture and expansion of the nasal floor. The TPD was activated for 1.5 mm at the completion of the osteotomy to facilitate separation of the midpalatal suture.




Fig. 2. Right nasal endoscopic view showing the midpalatal osteotomy.

2.3. *Expansion Process*

The TPD was activated between 5-7 days after surgery by 0.3 mm per day. The expansion process is deemed complete when either the patient has experienced no further clinical improvement with continual expansion or when there has been 7 mm expansion. Once expansion was completed, the TPD was locked and removed under local anesthesia two months later.

2.4. *Polysomnography*

PSG was performed within 2 years before surgery, and postoperative PSG was performed within 6 weeks after removal of the TPD. In-lab PSG included electroencephalography, electrooculogram, and electromyography of chin and leg movements; respirations were monitored with a nasal cannula, mouth thermistor, uncalibrated inductive plethysmography, thoracic and abdominal bands, snore microphone, position sensor and

finger pulse oximetry. PSG scoring was based on 2012 American Academy of Sleep Medicine (AASM) recommendations.

2.5. *Cone Beam Computed Tomography (CBCT)*

All patients underwent CBCT preoperatively and at 3 months postoperatively. CBCT scans were acquired in the supine position in extended field modus (FOV: 16x22cm, scanning time 2x20s, voxel size 0.4 mm, NewTom 3D VGI, Cefla North America, Charlotte, NC). Data from CBCT were exported in Digital Imaging and Communications in Medicine (DICOM) format. NNT software (QR sri, Verona, Italy) was used to viewing and OnDemand3D Fusion software (OnDemand3D Technology, Tustin, CA) was used for superimposition of pre- and post-treatment images for comparison and measurements. The measurement methods were derived from a previously published study (17).

2.6. *Questionnaires*

ESS and NOSE questionnaires were administered at preoperative appointments (1-3 weeks prior to surgery) and between 3-4 months postoperatively just prior to TPD removal.

2.7. *Statistical Analyses*

Descriptive statistics and frequency distributions were performed on demographic and clinical characteristics. Summary measures were computed as means and standard deviations for continuous variables or counts and proportions for categorical variables. The paired samples Wilcoxon signed-rank test was used to compare preoperative and postoperative parameters due to the small sample size and skewed distribution of some measures. Data were evaluated for extreme, implausible, and missing values. All analyses were performed using R Studio version 1.1.383 with a 2-sided p -value less than 0.05 to indicate statistical significance.

3. Results (Table 1)

Table 1. Preoperative and postoperative values for demographic and clinical characteristics of all patients (n = 33).

	Preoperative Mean ± SD or Count (%)	Postoperative Mean ± SD or Count (%)	p-value*
<i>Demographic Characteristics</i>			
Age (years)	29.4 ± 14.6	-	-
Male	18 (54.5%)	-	-
Female	15 (45.5%)	-	-
<i>Clinical Characteristics</i>			
Body Mass Index (BMI; kg/m²)	24.7 ± 2.6	25.0 ± 2.5	0.11
Anterior Nasal Spine (ANS) Expansion (mm)	-	4.9 ± 1.2	-
Posterior Nasal Spine (PNS) Expansion (mm)	-	5.6 ± 1.2	-
Dental Diastema Created (mm)	-	2.3 ± 0.8	-
Procedure Time (min)	-	54.0 ± 6.0	-
Oxygen Desaturation Index (ODI)	11.8 ± 9.6	1.8 ± 3.7	<0.0001
Apnea Hypopnea Index (AHI)	31.6 ± 11.3	10.1 ± 6.3	<0.0001
Minimum Oxygen Saturation (%)	89.4 ± 3.1	92.1 ± 2.1	<0.0001
Epworth Sleepiness Score (ESS)	13.4 ± 4.0	6.7 ± 3.1	<0.0001
Nasal Obstruction Septoplasty Effectiveness (NOSE)	57.8 ± 12.9	15.6 ± 5.7	<0.0001

*p-values determined using Wilcoxon signed-rank test

Thirty-three patients (18 males) were evaluated in this retrospective study. The mean age was 29.4±14.6 years (range 15-61). The AHI improved from 31.6±11.3 to 10.1±6.3, and the overall AHI reduction was 68%. The oxygen desaturation index (ODI) improved from 11.8±9.6 to 1.8±3.7 and the ESS score decreased from 13.4±4.0 to 6.7±3.1. The lowest oxygen saturation increased from 89.4±3.1 to 92.1±2.1. Nasal breathing improved as per the NOSE questionnaire from 57.8±12.9 to 15.6±5.7. The anterior nasal floor expanded 4.9±1.2 mm (measured at ANS), posterior nasal floor expanded 5.6±1.2 mm (measured at PNS), and the dental diastema created was 2.3±0.8 mm (range 1-5) at the completion of expansion (Figs. 3-5).




Fig. 3. (a) Preoperative palatal view. (b) Postoperative palatal view showing the TPD in place at the completion of expansion. (c) Preoperative frontal view. (d) Postoperative frontal view showing a 2 mm diastema despite 5 mm of nasal floor expansion (see CBCT showing the 5 mm widening in Figure 3d).




Fig. 4. CBCT of the patient in Figure 2. (a) Preoperative frontal view. (b) Postoperative frontal view at the completion of expansion showing a 5 mm opening at the ANS. (c) Preoperative palatal view. (d) Postoperative palatal view at the completion of expansion showing a 5 mm opening at the PNS. Note the small dental diastema.




Fig. 5. (a) Frontal view of CBCT of another patient at the initial phase of expansion with opening of the midpalatal suture. (b) Frontal view of CBCT showing continual expansion that resulted in greater opening of the nasal floor (5 mm separation of the ANS). (c) Palatal view of CBCT at the initial phase of expansion. Note the separation of the midpalatal suture. (d) Palatal view of CBCT demonstrating widening of the palate (PNS widened 6 mm). Note the absence of a dental diastema despite significant widening of the nasal floor.

The mean operative time was 54.0 ± 6.0 minutes. Body mass index (BMI) increased from 24.7 ± 2.6 to 25.0 ± 2.5 . Patients with mild to moderate OSA (19 patients) were discharged the same day, while patients with severe OSA were observed overnight and discharged the following morning. All patients returned to work and regular activities within 3 days.

31 patients (93.9%) experienced improvement and 29 patients (87.9%) noted significant improvement, which was defined as a > 50% reduction in AHI as well as reduction of ESS and NOSE scores. During and post-expansion, two of the patients who were bilevel-dependent noted improved symptoms while on bilevel with decreased AHI values, decreased IPAP and/or EPAP pressures, and reduced air leak (Fig. 6).

05/27/2018
Sunday night

CPAP Statistics
709 days of CPAP Data, between Fri Jun 17 2016 and Sun May 27 2018

Details	Most Recent	Last Week	Last 30 Days	Last 6 Months	Last Year
CPAP Usage					
Average Hours per Night	06:44	07:57	08:09	08:33	08:27
Compliance	100%	100%	100%	100%	100%
Therapy Efficacy					
AHI	0.30	1.19	1.01	2.04	72473.26
Obstructive Index	0.00	0.00	0.00	0.00	0.00
Hypopnea Index	0.30	1.15	0.99	1.97	2.40
Clear Airway Index	0.00	0.00	0.00	0.00	72470.78
Leak Statistics					
Average Leak Rate	0.61	4.99	4.54	15.89	22.53
90% Leak Rate	0.00	0.00	0.00	0.00	0.00
% of time above Leak Rate threshold	1.14%	5.87%	6.30%	20.29%	27.06%
Pressure Statistics					
Average EPAP	7.46	7.09	8.21	11.27	11.90
Min EPAP	4.06	4.06	4.06	4.06	4.06
Max EPAP	9.10	10.42	13.32	15.00	15.00
Average IPAP	11.75	11.17	12.50	16.71	17.33
90% IPAP	0.00	0.00	0.00	0.00	0.00
Min IPAP	7.02	4.80	4.20	4.20	4.20
Max IPAP	17.70	25.00	25.00	25.00	25.00

machine max

Fig. 6. Bilevel readings before and during the expansion process. Note the reduction in IPAP, EPAP, AHI and leak.

However, none of these patients were able to eliminate bilevel use and were thus considered surgical failures. Two other patients experienced improvement in nasal breathing as per improved NOSE scores, but these patients did not experience significant improvement in PSG or ESS scores and were thus also considered surgical failures. Despite insufficient clinical improvement, all four patients achieved successful maxillary expansion on CBCT.

Two patients experienced minor complications. Both patients misunderstood the instructions regarding expansion and had turned the expander in the opposite direction. Both patients required adjustment of the device under local anesthesia in the office setting.

4. Discussion

Maxillary expansion is a dental procedure that was originally designed to widen a narrow maxilla for the treatment of dental deformities. Numerous expanders have been used and include tooth-borne, bone-borne and hybrid (attached to teeth and bone) devices with or without SARPE. Although the objective of maxillary expansion is to expand bone rather than teeth, almost all of these expansion methods cause varying degrees of lateral tilting of the dentoalveolar component due to pressure from the expanders. When applying this maxillary expansion to treat OSA rather than to treat dental deformities, the goal is to expand the nasal floor as much as possible to enlarge the airway since expansion of the dentoalveolar component does not affect the airway at all. With the existing maxillary expansion procedures (excluding EASE), it is necessary to over-expand the dental alveolus to induce secondary widening of the nasal floor. This results in an undesired effect in which the degree of dental widening greatly exceeds the degree of nasal floor widening and results in an unsightly, large diastema and significant malocclusion. Such traditional approaches to maxillary expansion increase the orthodontic treatment time and risks jeopardizing the vitality of teeth (18, 19). More importantly, the extent of nasal floor widening is inconsistent, inadequate and often nonexistent (20-22), especially in the posterior nasal region. This is because the posterior region of the maxilla has been shown to be the most resistant to expansion (9, 22). Traditional maxillary expansion procedures thus cause an undesirable fan shape expansion characterized by excessive anterior dental widening and minimal to no posterior nasal airway expansion. Although this expansion pattern may improve nasal breathing, we have found that this expansion pattern does not significantly improve OSA (unpublished data). We postulate that the posterior nasal floor is the most important area to expand when optimizing airway size to treat OSA. Widening of the posterior hard palate not only increases nasal airway volume, but also may expand the retropalatal airway region because several palatopharyngeal muscles originate from the posterior hard palate.

The results of this study demonstrated that EASE is a novel surgical approach that can be used to achieve consistent expansion of the nasal floor throughout the entire nasal region to treat OSA. Interestingly, in contrast to traditional maxillary expansion procedures (18, 23, 24), EASE not only achieved the greatest degree of airway expansion in the posterior

nasal floor, but also did so with a much smaller resultant anterior diastema. We believe the use of a bone-borne expander applied in posterior maxilla, along with strategic osteotomy at a high-stress region of the skull, enables EASE to apply adequate forces to separate the midpalatal sutures and expand the nasal floor while preventing tilting of the teeth. Strategic separation of the midpalatal suture, including separation of the PNS, and strategic separation of the pterygomaxillary suture are essential in achieving adequate nasal floor separation. Indeed, the importance of pterygomaxillary separation to facilitate posterior maxillary widening was previously reported with the use of TPD and traditional SARPE technique (25). Clearly, a less invasive surgical approach in optimizing airway improvement while minimizing dental changes to reduce postoperative risks and orthodontic treatment time are preferred.

The nasal endoscopic surgical approach (EASE) is considerably less invasive compared to other methods because EASE obviates the need for incisions or bone cuts adjacent to the upper lip, gums, sinus walls and teeth. EASE thus prevents any long-term distortion of the nose and/or upper lip that may have otherwise resulted from muscle stripping and trauma that occur in other surgical maxillary expansion procedures. The avoidance of traditional osteotomy near the tooth roots reduces postoperative pain and swelling while preventing injury to the teeth and bone. The use of piezoelectric blade simultaneously cauterizes the nasal mucosal, thus reducing the risk of bleeding. EASE also enables patients to use PAP therapy for airway protection immediately following surgery. With other methods of maxillary expansion that include incisions and osteotomies at the anterior maxilla and sinus walls, PAP therapy cannot be used immediately postoperatively due to the risk of subcutaneous emphysema.

The majority of the patients in this study experienced improvement in nasal breathing and OSA symptoms within the first week of expansion. Of note, two of the patients who were bilevel-compliant prior to surgery continued bilevel therapy immediately after surgery. The bilevel pressure had to be reduced shortly after initiation of the expansion. This is quite interesting and suggests that even a slight widening of the entire nasal airway had a significant impact on airway patency.

Many factors influence a patient's decision to have surgery to treat his or her OSA. The invasiveness of the operation, extent of improvement, recovery time and time off from work are major considerations. The minimally-invasive nature, low complication rate and short operative duration characteristic of EASE allowed patients to be discharged the same day and return to school or work within a few days. Such benefits of EASE do not exist with other known maxillary expansion surgical techniques.

5. Conclusions

The results of this study demonstrate that OSA and nasal breathing can be improved via EASE of the nasal floor in adolescents and adults. As a novel and minimally-invasive maxillary widening surgical procedure, EASE allows patients to return quickly to their daily activities and has numerous additional advantages compared to that of other surgical widening methods.

Funding Sources

This study was not funded.

Conflicts of Interest

The authors had no conflicts of interest to declare in relation to this article.

References

- (1) Cistulli PA, Palmisano RG, Poole MD. Treatment of obstructive sleep apnea syndrome by rapid maxillary expansion. *Sleep* 1998;21:831-5.
- (2) Pirelli P, Saponara M, Guilleminault C. Rapid maxillary expansion in children with obstructive sleep apnea syndrome. *Sleep* 2004;27:761-6.
- (3) Quo SD, Hyunh N, Guilleminault C. Bimaxillary expansion therapy for pediatric sleep-disordered breathing. *Sleep Medicine* 30;45-51, 2017.
- (4) Camacho M, Chang ET, Song SA, et al. Rapid maxillary expansion for pediatric obstructive sleep apnea: a systematic review and meta-analysis. *Laryngoscope* 2017;127:1712-9.

- (5) Villa MP, Rizzoli A, Miano S, et al. Efficacy of rapid maxillary expansion in children with obstructive sleep apnea syndrome: 36 months of follow-up. *Sleep Breath* 2011;15:179-184.
- (6) Persson M, Thilander B. Palatal suture closure in man from 15-35 years of age. *Am J Orthod* 1977;72:42-52.
- (7) Melsen B, Melsen F. The postnatal development of the palatomaxillary region studied on human autopsy material. *Am J Orthod* 1982;82:329-342.
- (8) Koudstaal MJ, Poort LJ, van der Wal KG, et al. Surgically assisted rapid maxillary expansion: a review of the literature. *Int J Oral Maxillofac Surg* 2005;34:709-714.
- (9) Shetty V, Caridad JM, Caputo AA, et al. Biomechanical rationale for surgical-orthodontic expansion of the adult maxilla. *J Oral Maxillofac Surg* 1994;52:742-9.
- (10) Kurt G, Altug AT, Turker G, et al. Effects of surgical and nonsurgical rapid maxillary expansion on palatal structures. *J Craniofac Surg* 2017;28:775-780.
- (11) Dergin G, Aktop S, Varol A, et al. Complications related to surgically assisted rapid palatal expansion. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2015;119:601-7.
- (12) Williams BJ, Currimbhoy S, Silva A, et al. Complications following surgically assisted rapid palatal expansion: a retrospective cohort study. *J Oral Maxillofac Surg* 2012;70:2394-2402.
- (13) Herford AS, Akin L, Cicciu M. Maxillary vestibular incision for surgically assisted rapid palatal expansion: evidence for a conservative approach. *Orthodontics* 2012;13:168-75.
- (14) Berger JL, Pangrazio-Kulbersh V, Thomas BW, et al. Radiographic analysis of facial changes associated with maxillary expansion. *Am J Orthod Dentofacial Orthop* 1999;116:563-71.
- (15) Ferrario VF, Sforza C, Schmitz JH, et al. Three-dimensional facial morphometric assessment of soft tissue changes after orthognathic surgery. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1999;88:549-56.
- (16) Guilleminault C, Li KK. Maxillomandibular expansion for the treatment of sleep-disordered breathing: preliminary result. *Laryngoscope* 2004;114:893-6.

- (17) Nada RM, van Loon B, Schols JG, et al. Volumetric changes of the nose and nasal airway 2 years after tooth-borne and bone-borne surgically assisted rapid maxillary expansion. *Eur J Oral Sci* 2013;121:450-456.
- (18) Liu S, Guilleminault C, Huon LK, et al. Distraction osteogenesis maxillary expansion for adult obstructive sleep apnea patients with high arched palate. *Oto Head Neck Surg* 2017;157:345-348.
- (19) Pereira MD, Koga AF, Prado GP, et al. Complications from surgically assisted rapid maxillary expansion with Haas and Hyrax expanders. *J Craniofac Surg* 2018;29:275-278.
- (20) Deeb W, Hansen L, Hotan T, et al. Changes in nasal volume after surgically assisted bone-borne rapid maxillary expansion. *Am J Orthod Dentofacial Orthop* 2010;137:782-9.
- (21) Pereira MD, Prado GP, Abramoff MM, et al. Classification of midpalatal suture opening after surgically assisted rapid maxillary expansion using computed tomography. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010;110:41-5.
- (22) De Assis DS, Xavier TA, Noritomi PY, et al. Finite element analysis of bone stress after SARPE. *J Oral Maxillofac Surg* 2014;72:167.e1-7.
- (23) Asscherickx K, Govaerts E, Aerts J, Vande Vannet B. Maxillary changes with bone-borne surgically assisted rapid palatal expansion: a prospective study. *Am J Orthod Dentofacial Orthop* 2016;149:374-83.
- (24) Park JJ, Park YC, Lee KJ, et al. Skeletal and dentoalveolar changes after miniscrew-assisted rapid palatal expansion in young adults: a cone-beam computed tomography study. *Korean J Orthod* 2017;47:77-86.
- (25) Matterini C, Mommaerts MY. Posterior transpalatal distraction with pterygoid disjunction: a short-term model study. *Am J Orthod Dentofacial Orthop* 2001;120:498-502.

- Endoscopically-assisted surgical expansion widens the entire nasal floor
- Endoscopically-assisted surgical expansion improves obstructive sleep apnea
- Endoscopically-assisted surgical expansion improves nasal breathing